

  
XPOVIO<sup>®</sup>  
(selinexor)<sup>20 mg</sup>  
tablet

## Getting started with XPOVIO<sup>®</sup> (selinexor)

the first and only FDA-approved  
oral medicine of its kind

XPOVIO is a prescription medicine used to treat adults with certain types of diffuse large B-cell lymphoma (DLBCL) that has come back (relapsed) or that did not respond to previous treatment (refractory) and who have received at least 2 prior therapies.

It is not known if XPOVIO is safe and effective in children less than 18 years of age.

### What is the most important information I should know about XPOVIO?

**Blood Tests:** Your healthcare provider will do blood tests before you start taking XPOVIO, and often during the first 3 months of treatment, and then as needed during treatment to monitor you for side effects. Your healthcare provider may change your dose of XPOVIO, stop your treatment for a period of time, or completely stop your treatment based on tests performed during treatment with XPOVIO.

**Tell your healthcare provider right away if you have any bleeding or easy bruising during treatment with XPOVIO.**

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).





### WHAT IS XPOVIO?

XPOVIO® (selinexor) is a prescription medicine used to treat adults with certain types of diffuse large B-cell lymphoma (DLBCL) that has come back (relapsed) or that did not respond to previous treatment (refractory) and who have received at least 2 prior therapies.

It is not known if XPOVIO is safe and effective in children less than 18 years of age.

### What is the most important information I should know about XPOVIO? (continued)

- **Low platelet counts.** Low platelet counts are common with XPOVIO and can lead to bleeding, which can be severe and can sometimes cause death. Your healthcare provider may prescribe platelet transfusions or other treatments for your low platelet counts.
- **Low white blood cell counts.** Low white blood cell counts are common with XPOVIO and can sometimes be severe. You may have an increased risk of getting bacterial infections during treatment with XPOVIO. If needed, your healthcare provider may prescribe antibiotics if you have signs or symptoms of infection, or may prescribe certain medicines to help increase your white blood cell count.

 **XPOVIO**<sup>®</sup>  
(selinexor) <sup>20 mg</sup> tablet

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).

# Table of contents

---

Why XPOVIO® (selinexor)? .....	4
What is XPO1 and why does it matter? .....	6
How does XPOVIO work to block XPO1? .....	7
How did patients respond to treatment with XPOVIO? .....	8
How long did responses to XPOVIO last? .....	9
What are the possible side effects with XPOVIO? .....	10
How do I take XPOVIO? .....	11
What else should I know about taking XPOVIO? .....	12
What should I ask my healthcare team before taking XPOVIO? .....	13
<b>For caregivers:</b> What should I know about caring for someone taking XPOVIO? .....	14
What support and resources are available? .....	15

## What is the most important information I should know about XPOVIO? (continued)

- **Nausea and vomiting and/or diarrhea.** Nausea and vomiting and/or diarrhea can occur when you take XPOVIO and can sometimes be severe. You may be at risk for becoming dehydrated and may need treatments. Your healthcare provider may prescribe anti-nausea or anti-diarrhea medicines for you to take before you start and during treatment with XPOVIO.

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).





## Why XPOVIO® (selinexor)?

**XPOVIO** is an FDA-approved medicine that treats adults with certain types of diffuse large B-cell lymphoma (DLBCL) that has come back (relapsed) or that did not respond to previous treatment (refractory) and who have received at least 2 prior therapies.

### What is the most important information I should know about XPOVIO? (continued)

- **Loss of appetite and weight loss.** Loss of appetite and weight loss are common with XPOVIO and can sometimes be severe. Tell your healthcare provider if you have a decrease or loss of appetite and if you are losing weight. Your healthcare provider may prescribe medicines or advise on nutritional support.

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).

**XPOVIO**<sup>®</sup>  
(selinexor) 20 mg  
tablet

---

## XPOVIO® (selinexor) is the **only FDA-approved oral medicine of its kind**

---



**Taken by mouth**  
at home



Does **not** need to be taken  
with any other anti-cancer  
medications for DLBCL



May work **when other**  
**treatments don't**



**Works at the core**  
**of the cell** (the nucleus)  
to kill some cancer cells

### What is the most important information I should know about XPOVIO? (continued)


- **Decreased sodium levels in your blood.** Decreased sodium levels in your blood are common with XPOVIO and can also sometimes be severe. You may not have any symptoms. Your healthcare provider may talk with you about your diet and prescribe IV fluids or salt tablets for you.

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).

  
XPOVIO®  
(selinexor) 20 mg  
tablet

# What is XPO1 and why does it matter?

---




*For illustrative purposes only.*

## Healthy cells


- Healthy cells have a protein called XPO1 that is responsible for carrying important materials in and out of the core of the cell (the nucleus)
- Many of these materials help fight cancer, but only when they are inside the nucleus

# How does XPOVIO® (selinexor) work to block XPO1?

Some cancer cells have too much XPO1



In this cancer cell not exposed to XPOVIO




XPO1 takes too much of the important materials that fight cancer out of the nucleus.

cancer cell growth and survival

*For illustrative purposes only.*

In this cancer cell exposed to XPOVIO



**XPOVIO targets and blocks XPO1**

XPO1 can no longer carry the important anti-cancer materials out of the nucleus. These materials accumulate in the nucleus.

cancer cell death

*For illustrative purposes only.*

## What is the most important information I should know about XPOVIO? (continued)


- **Serious infections.** Infections can occur when you are taking XPOVIO and can be serious and can sometimes cause death. This includes upper or lower respiratory tract infections, such as pneumonia, and an infection throughout your body (sepsis). **Tell your healthcare provider right away if you have any signs or symptoms of an infection such as cough, chills, or fever during treatment with XPOVIO.**

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).

**XPOVIO**<sup>®</sup>  
(selinexor) 20 mg tablet

## How did patients respond to treatment with XPOVIO® (selinexor)?

In the clinical trial, 134 adults with relapsed or refractory DLBCL (RR DLBCL) were treated with XPOVIO 60 mg on Days 1 and 3 of every week. FDA approval was based upon the results of this trial, and additional trials may be needed.



**29%**

**of patients responded to treatment with XPOVIO**


### What is the most important information I should know about XPOVIO? (continued)

- **Neurologic side effects.** XPOVIO can cause dizziness, fainting, decreased alertness, and changes in your mental status, including confusion and decreased awareness of things around you. XPOVIO may also cause problems with thinking (cognitive problems), seeing or hearing things that are not really there (hallucinations), and may cause you to become very sleepy or drowsy. These problems can sometimes be severe and life-threatening. Tell your healthcare provider right away if you get any of these symptoms.

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).

**XPOVIO**<sup>®</sup>  
(selinexor) 20 mg  
tablet

# How long did responses to XPOVIO® (selinexor) last?



The median duration of response (DOR) was a pre-specified secondary outcome, determined by a particular statistical analysis. Limitations to this analysis for the data from this study include a small number of patients and the effect of data from patients with prolonged remission durations. Approximately half of the responders in this analysis had a DOR lasting less than 3 months. The analysis of DOR is presented alongside an analysis demonstrating the number of patients responding at certain timepoints (3, 6, 12 months).

The **median** number is the middle point of a set of numbers, in which half the numbers are above the median and half are below.

## What is the most important information I should know about XPOVIO? (continued)

- **Do not drive or operate heavy or dangerous machinery** until you know how XPOVIO affects you. XPOVIO may cause dizziness and confusion. Take precautions to prevent a fall.

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).

**XPOVIO**<sup>®</sup>  
(selinexor)<sup>20 mg</sup> tablet

## What are the possible side effects with XPOVIO® (selinexor)?

### Serious side effects

- Low platelet counts
- Low white blood cell counts
- Nausea and vomiting
- Diarrhea
- Loss of appetite and weight loss
- Decreased sodium levels in your blood
- Infections
- Neurologic side effects

These are not all the possible side effects of XPOVIO.

**Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088 or [fda.gov/medwatch](https://www.fda.gov/medwatch).**

### Common side effects

- Tiredness
- Low red blood cell count (anemia). Symptoms may include tiredness and shortness of breath
- Constipation
- Shortness of breath
- Increased blood sugar
- Changes in body salt and mineral levels in your blood
- Changes in kidney and liver function blood tests

 XPOVIO®  
(selinexor) 20 mg tablet

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).

# How do I take XPOVIO® (selinexor)?

The recommended dosage of XPOVIO is 60 mg taken by mouth on Days 1 and 3 of each week



- Days 1 and 3 depend on which day of the week you start XPOVIO. For example, if you start XPOVIO on Tuesday, Tuesday is Day 1 and Thursday is Day 3
- Your healthcare provider may prescribe medicines to help prevent nausea/vomiting during treatment with XPOVIO
- Take your medication as prescribed by your healthcare provider and be sure to let them know how you are feeling along the way

**During treatment, your healthcare provider may change your dose of XPOVIO. Always take XPOVIO exactly as prescribed.**

## What is the most important information I should know about XPOVIO? (continued)

### Common side effects of XPOVIO include:

- tiredness
- low red blood cell count (anemia). Symptoms may include tiredness and shortness of breath
- constipation
- shortness of breath
- increased blood sugar
- changes in body salt and mineral levels in your blood
- changes in kidney and liver function blood tests

These are not all of the possible side effects of XPOVIO.

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).

  
**XPOVIO**<sup>®</sup>  
(selinexor) 20 mg  
tablet

## What else should I know about taking XPOVIO® (selinexor)?

---

- Talk to your healthcare provider before you change your dose or stop taking XPOVIO
- Swallow XPOVIO whole with water. Never break, chew, crush, or divide the tablet
- Take any medicines your healthcare provider prescribes for nausea/vomiting before and during treatment with XPOVIO
- Tell your healthcare provider if the medicines don't help with nausea/vomiting
- Drink enough fluids during treatment to prevent dehydration
- Eat enough calories to prevent weight loss during treatment
- If you miss a dose of XPOVIO, take your next dose at your next regularly scheduled day and time
- If you vomit after taking a dose of XPOVIO, do not take an extra dose. Take your next dose at your next regularly scheduled day and time
- If you take too much XPOVIO, call your healthcare provider right away
- Do not share your XPOVIO prescription with other people
- Your healthcare provider will do blood tests before you start taking XPOVIO, more often during the first 3 months of treatment, and then as needed

### What is the most important information I should know about XPOVIO? (continued)

**Before taking XPOVIO, tell your healthcare provider about all of your medical conditions, including if you:**

- have or have had a recent or active infection
- have or have had bleeding problems
- are pregnant or plan to become pregnant. XPOVIO can harm your unborn baby
- are taking prescription and over-the-counter medicines, vitamins, and herbal supplements

**Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).**



## What should I ask my healthcare team before taking XPOVIO® (selinexor)?

- Which **anti-nausea medicines** will I take prior to treatment with XPOVIO?
- How often will I need a **blood test** while taking XPOVIO?
- How do I ensure I **stay hydrated** while taking XPOVIO?
- Should I expect to **lose weight** while taking XPOVIO? How much?
- Will my **dose** be changed throughout treatment with XPOVIO?
- How will my **side effects** be managed?
- What should I do if I have a **new side effect or symptom**?
- What should I know if I'm taking **other medications**?

**Always ask your healthcare provider any questions you may have about treatment with XPOVIO.**

### What is the most important information I should know about XPOVIO? (continued)

**Ability to have children:** XPOVIO may affect the ability of both women and men to have children. Talk to your healthcare provider if you have concerns about fertility.

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).

 **XPOVIO**<sup>®</sup>  
(selinexor) 20 mg tablet



FOR CAREGIVERS:

## What should I know about caring for someone taking XPOVIO® (selinexor)?

### When caring for patients taking XPOVIO, keep in mind:

- The **dosing schedule** for XPOVIO and anti-nausea medicines will be determined by the healthcare provider
- **Call the healthcare provider** if your loved one experiences any side effects
- The healthcare team will do **blood tests** before your loved one starts taking XPOVIO, more often during the first 3 months of treatment, and then as needed
- Encourage your loved one to **drink lots of fluids and eat nutritious food** while on XPOVIO

### What is the most important information I should know about XPOVIO? (continued)

**Females who are able to become pregnant:** Your healthcare provider will check to see if you are pregnant before you start taking XPOVIO. You should use effective birth control (contraception) during treatment with XPOVIO and for 1 week after your last dose, as XPOVIO can harm an unborn baby. Tell your healthcare provider right away if you become pregnant or think you might be pregnant during treatment with XPOVIO. Do not breastfeed during treatment with XPOVIO and for 1 week after your last dose of XPOVIO. It is not known if XPOVIO passes into your breast milk.

**Males with female partners who are able to become pregnant** should use effective birth control during treatment with XPOVIO and for 1 week after your last dose.

To report **SUSPECTED ADVERSE REACTIONS**, contact Karyopharm Therapeutics Inc. at 1-888-209-9326 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).

 **XPOVIO**<sup>®</sup>  
(selinexor) 20 mg  
tablet

# What support and resources are available?

KaryForward is a patient support program by Karyopharm Therapeutics dedicated to providing assistance and resources to you and your caregiver for XPOVIO® (selinexor)

The KaryForward program is here to help you every step of the way during your treatment journey:



## Insurance coverage\*

- Understand your benefits
- File claims
- Provide support during insurance appeals
- Find out what is covered by your insurance
- Access Karyopharm medications, even if there are unexpected delays or interruptions



## Financial assistance

- Qualify for the XPOVIO Copay Program and pay as little as \$5 for each XPOVIO prescription, or
- If eligible, get XPOVIO at no cost to you



## Support and resources

- Nurse Case Managers who can provide education and support during treatment
- Referrals to additional third-party support, such as transportation assistance



**IF YOU'RE READY TO ENROLL  
OR HAVE QUESTIONS ABOUT  
KARYFORWARD:**

**CALL US**

**1-877-KARY4WD (1-877-527-9493)**  
Monday through Friday, 8 AM to 8 PM ET

**VISIT US**

**KaryForward.com**

\*Reimbursement Specialists are available to answer questions about the insurance approval process.

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).

**XPOVIO®**  
(selinexor) 20 mg  
tablet

# XPOVIO® (selinexor) is the first and only FDA-approved oral medicine of its kind

- **Works at the core of the cell** (the nucleus) to kill some cancer cells
- **May work when other treatments don't**
- XPOVIO is **taken by mouth** at home

The most serious side effects with XPOVIO include low platelet counts, low white blood cell counts, nausea and vomiting, diarrhea, loss of appetite and weight loss, decreased sodium levels in your blood, infections, and neurologic side effects. Common side effects include tiredness, low red blood cell count (anemia), constipation, shortness of breath, increased blood sugar, changes in body salt and mineral levels in your blood, and changes in kidney and liver function blood tests.

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Medication Guide](#).



For more information, call 1-888-209-9326.  
© 2020 Karyopharm Therapeutics Inc. All rights reserved. US-XPOV-03/20-00036



Learn more at [XPOVIO.com](https://www.xpovio.com)

**GET ASSISTANCE AND RESOURCES FOR XPOVIO TREATMENT WITH THE KARYFORWARD® PROGRAM.**

CALL

**1-877-KARY4WD (1-877-527-9493)**  
Monday through Friday, 8 AM to 8 PM ET

VISIT

[KaryForward.com](https://www.KaryForward.com)

 **XPOVIO®**  
(selinexor) 20 mg tablet