

Understanding Follicular Lymphoma: Relapsed/Refractory

Follicular lymphoma (FL) is the most common *indolent* (slow-growing) form of non-Hodgkin Lymphoma (NHL), accounting for just over 10 percent of all B-cell NHLs.

Common symptoms of FL include swelling of the lymph nodes in the neck, underarms, abdomen, or groin; fatigue; and much less commonly, fevers, night sweats, and weight loss. Often, patients with FL have no obvious symptoms of the disease and the diagnosis is found incidentally during a routine annual checkup or following imaging studies for unrelated reasons. Many patients may not initially need treatment for some time.

FL is generally very responsive to radiation and chemotherapy and many patients go into a *remission* (disappearance of signs and symptoms) that lasts for years after their initial treatment; however, the disease often returns. For patients who *relapse* (disease returns after treatment) or become *refractory* (disease no longer responds to treatment), *second-line therapies* (treatment given when initial therapy does not work or stops working) are often successful in providing another remission. Some patients who relapse do not need treatment right away, and an *active surveillance* (also known as “watchful waiting” or “careful observation”) approach might be used. With this strategy, patients’ overall health and disease are monitored through regular physical exams and sometimes periodic imaging tests. Active treatment is started if the patient begins to develop lymphoma-related symptoms or there are signs that the disease is progressing based on testing during follow-up visits. For those who need treatment, the same therapies used for newly diagnosed patients can often be used in patients with relapsed/refractory FL, but additional treatments are also available.



TREATMENT OPTIONS

Treatment for relapsed/refractory FL is based on a patient’s age, overall health, symptoms, prior therapy or therapies, and the duration of remission from the last treatment they received. Chemotherapy, radiation, and monoclonal antibodies such as rituximab (Rituxan [for intravenous infusion] and Rituxan Hycela [for subcutaneous injection after the first IV infusion of rituxan]) and obinutuzumab (Gazyva) may be used to treat relapsed/refractory FL.

Common second-line regimens include:

- Bendamustine (Treanda) with or without rituximab (Rituxan), if not used for first-line treatment
- Copanlisib (Aliqopa)
- Duvelisib (Copiktra)
- Idelalisib (Zydelig)
- Lenalidomide (Revlimid) with or without rituximab (often referred to as R2 [R-squared] when used in combination with rituximab)
- Obinutuzumab (Gazyva) with bendamustine, if bendamustine was not used in first-line treatment
- R-CHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisone)

- R-CVP (rituximab, cyclophosphamide, vincristine, and prednisone)
- Rituximab alone
- Tazemetostat (approved for relapsed/refractory FL with an EZH2 mutation or for patients with relapsed/refractory FL who have no satisfactory alternative treatment option)

Although uncommon, radioimmunotherapy (RIT) using an agent such as yttrium-90 ibritumomab tiuxetan (Zevalin), which is a radioactive particle connected to an antibody that targets cancer cells, may also be used alone or in combination with chemotherapy to treat relapsed/refractory FL. To learn more about RIT, view the *Radioimmunotherapy* fact sheet on the Lymphoma Research Foundation’s (LRF’s) website at lymphoma.org/publications.

Radiation therapy can be effective in some patients with relapsed/refractory FL who have localized disease. Often very low doses of radiation can be quite beneficial.

For some patients with relapsed/refractory FL, high-dose chemotherapy followed by stem cell transplantation may be an option. For more information on transplantation, view the *Understanding the Stem Cell Transplantation Process* publication on LRF’s website at lymphoma.org/publications.

RESPONSE TO RETREATMENT

With newer therapeutic regimens, many patients can achieve remissions after second-line or third-line treatments. Remissions are frequently shorter with each round of therapy, however remissions in the range of one year and longer can be seen with some of the treatments.

TRANSFORMED FL

Some patients with FL may eventually develop a transformed lymphoma, which occurs at a rate of three percent per year and is often more aggressive and usually requires more intensive types of treatment similar to aggressive lymphomas. The risk of developing a transformed lymphoma increases each year from the time of diagnosis until approximately 10 years afterwards, after which point transformations become rare. For more information on transformed lymphomas, view the *Transformed Lymphomas* fact sheet on LRF's website at lymphoma.org/publications.

TREATMENTS UNDER INVESTIGATION

Many treatments are currently being studied in clinical trials alone or as part of a combination therapy regimen in patients with relapsed/refractory FL. Some of these treatments include:

- Acalabrutinib (Calquence)
- ALT-803
- ASN002
- Atezolizumab (Tecentriq)
- Axicabtagene ciloleucel (Yescarta)
- BGB-3111
- Blinatumomab (Blincyto)
- Buparlisib
- Cerdulatinib
- Denintuzumab mafodotin
- Durvalumab (Imfinzi)
- Entospletinib
- Ibrutinib (Imbruvica)
- Idasanutlin
- INCB050465
- Ixazomib (Ninlaro)
- JCAR017
- Nivolumab (Opdivo)
- Pembrolizumab (Keytruda)
- PLX2853
- Romidepsin (Istodax)

- SAIT101
- SD-101
- Tisagenlecleucel (Kymriah)
- Ublituximab
- Umbralisib
- Vecabrutinib
- Venetoclax (Venclexta)

It is critical to remember that today's scientific research is continuously evolving. Treatment options may change as new treatments are discovered and current treatments are improved. Therefore, it is important that patients check with their physician or with LRF for any treatment updates that may have recently emerged.

CLINICAL TRIALS

Clinical trials are crucial in identifying effective drugs and determining optimal doses for patients with lymphoma. Patients interested in participating in a clinical trial should view the *Understanding Clinical Trials* fact sheet on LRF's website at lymphoma.org/publications and the *Clinical Trials Search Request Form* at lymphoma.org, talk to their physician, or contact the LRF Helpline for an individualized clinical trial search by calling (800) 500-9976 or emailing helpline@lymphoma.org.

FOLLOW-UP

Since FL is generally characterized by multiple disease relapses after responses to a variety of treatments, patients should have regular visits with a physician who is familiar with their medical history and the treatments they have received. Medical tests (such as blood tests, computed tomography [CT] scans, and positron emission tomography [PET] scans) may be required at various times during remission to evaluate the need for additional treatment.

Some treatments can cause long-term side effects or late side effects, which can vary based on duration and frequency of treatments, age, gender, and the overall health of each patient at the time of treatment. A physician will check for these side effects during follow-up care. Visits may become less frequent the longer the disease remains in remission.

Patients and their caregivers are encouraged to keep copies of all medical records and test results as well as information on the types, amounts, and duration of all treatments received. This documentation will be important for keeping track of any side effects resulting from treatment or potential disease recurrences. LRF's award-winning *Focus On Lymphoma* mobile app (lymphoma.org/mobileapp) can help patients manage this documentation.



PATIENT AND CAREGIVER SUPPORT SERVICES

A lymphoma diagnosis often triggers a range of feelings and concerns. In addition, cancer treatment can cause physical discomfort. One-to-one peer support programs, such as LRF's *Lymphoma Support Network*, connect patients and caregivers with volunteers who have experience with lymphoma or FL and similar treatments or challenges, for mutual emotional support and encouragement. Patients and loved ones may find this information useful whether the patient is newly diagnosed, in treatment, or in remission.

Resources

LRF offers a wide range of resources that address treatment options, the latest research advances, and ways to cope with all aspects of FL, including our award-winning mobile app. LRF also provides many educational activities, from in-person meetings to webinars, for people with lymphoma, as well as FL e-Updates that provide the latest disease-specific news and treatment options. To learn more about any of these resources, visit our websites at lymphoma.org/FL or lymphoma.org, or contact the LRF Helpline at (800) 500-9976 or helpline@lymphoma.org.

Contact LRF:

Helpline: (800) 500-9976

Email: helpline@lymphoma.org

www.lymphoma.org

Medical reviewer:

Isidore S. Lossos, MD

Sylvester Comprehensive Cancer Center

Supported through grants from:

Genentech
A Member of the Roche Group

Biogen

Bristol Myers Squibb

Epizyme

The *Understanding Lymphoma* series is published by the Lymphoma Research Foundation (LRF) for the purpose of informing and educating readers. Facts and statistics were obtained using published information, including data from the Surveillance, Epidemiology, and End Results (SEER) Program. Because each person's body and response to treatment is different, no individual should self-diagnose or embark upon any course of medical treatment without first consulting with his or her physician. The medical reviewer, the medical reviewer's institution, and LRF are not responsible for the medical care or treatment of any individual.

© 2020 Lymphoma Research Foundation

Last updated 2020